

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.



Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

PCT/US 96/02957

BAR CODE LABEL 		U.S. PATENT APPLICATION			
SERIAL NUMBER 08/397,651		FILING DATE 03/01/95	CLASS 435	GROUP ART UNIT 1802	
APPLICANT	ULF ERIKSSON, BALSTA, SWEDEN; BIRGITTA OLOFSSON, SUNDBYBERG, SWEDEN.				
	CONTINUING DATA*** VERIFIED _____				
	FOREIGN/PCT APPLICATIONS*** VERIFIED _____				
<div style="border: 2px solid black; padding: 5px; transform: rotate(-5deg); display: inline-block;"> PRIORITY DOCUMENT </div>					
STATE OR COUNTRY SEK	SHEETS DRAWING 3	TOTAL CLAIMS 23	INDEPENDENT CLAIMS 2	FILING FEE RECEIVED \$926.00	ATTORNEY DOCKET NO. 1064/41979
ADDRESS	EVENSON MCKEOWN EDWARDS AND LEMAHAN 1200 G STREET NW SUITE 700 WASHINGTON DC 20005				
TITLE	VASCULAR ENDOTHELIAL GROWTH FACTOR-B AND DNA CODING THEREFOR				
This is to certify that annexed hereto is a true copy from the records of the United States Patent and Trademark Office of the application which is identified above. By authority of the COMMISSIONER OF PATENTS AND TRADEMARKS Date APR 18 1996 Certifying Officer 					

08 392851

Inventors: Ulf Eriksson
Birgitta Olofsson

VASCULAR ENDOTHELIAL GROWTH FACTOR-B AND DNA CODING THEREFOR

Background of the Invention

Angiogenesis, or the proliferation of new capillary blood vessels, is a fundamental process necessary for normal growth and development of tissues. It is a prerequisite for the development and differentiation of the vascular tree, as well as for a wide variety of fundamental physiological processes including embryogenesis, somatic growth, tissue and organ repair and regeneration, cyclical growth of the corpus luteum and endometrium, and development and differentiation of the nervous system. In the female reproductive system, angiogenesis occurs in the follicle during its development, in the corpus luteum following ovulation and in the placenta to establish and maintain pregnancy. Angiogenesis additionally occurs as part of the body's repair processes, e.g. in the healing of wounds and fractures. Angiogenesis is also a factor in tumor growth, since a tumor must continuously stimulate growth of new capillary blood vessels in order to grow.

Capillary blood vessels consist of endothelial cells and pericytes. These two cell types carry all of the genetic information to form tubes, branches and entire capillary networks. Specific angiogenic molecules can initiate this process. In view of the physiological importance of angiogenesis, much effort has been devoted to the isolation, characterization and purification of factors that can stimulate angiogenesis, and a number of polypeptides which stimulate angiogenesis have been purified and characterized as to their molecular, biochemical and

biological properties. For reviews of such angiogenesis regulators, see Klagsbrun et al., "Regulators of Angiogenesis", *Ann. Rev. Physiol.*, 53:217-39 (1991); and Folkman et al., "Angiogenesis," *J. Biol. Chem.*, 267:10931-934 (1992).

One such growth factor, which is highly specific as a mitogen for vascular endothelial cells, is termed vascular endothelial growth factor (VEGF). See Ferrara et al., "The Vascular Endothelial Growth Factor Family of Polypeptides," *J. Cellular Biochem.*, 47:211-218 (1991); Connolly, "Vascular Permeability Factor: A Unique Regulator of Blood Vessel Function," *J. Cellular Biochem.*, 47:219-223 (1991). VEGF is a potent vasoactive protein that has been detected in media conditioned by a number of cell lines including bovine pituitary follicular cells. VEGF is a glycosylated cationic 46-48 kD dimer made up of two 24 kD subunits. It is inactivated by sulfhydryl reducing agents, resistant to acidic pH and to heating, and binds to immobilized heparin. VEGF is sometimes referred to as vascular permeability factor (VPF) because it increases fluid leakage from blood vessels following intradermal injection. It also has been called by the name vasculotropin.

Four different molecular species of VEGF have been detected. The 165 amino acid species has a molecular weight of approximately 46 kD and is the predominant molecular form found in normal cells and tissues. A less abundant, shorter form with a deletion of 44 amino acids between positions 116 and 159 (VEGF₁₂₁), a longer form with an insertion of 24 highly basic residues in position 116 (VEGF₁₆₅), and another longer form with an insertion of 41 amino acids (VEGF₁₈₉), which includes the 24 amino acid insertion found in VEGF₁₆₅, are also known. VEGF₁₂₁ and VEGF₁₆₅ are soluble proteins. VEGF₁₂₁ and VEGF₁₆₅ appear to be mostly cell-associated. All of the versions of VEGF are biologically active. For example, each of the species when applied intradermally is able to induce extravasation of Evans blue.

The various species of VEGF are encoded by the same gene and arise from alternative splicing of messenger RNA. This conclusion is supported by Southern blot analysis of human genomic DNA, which shows that the restriction pattern is identical using either a probe for VEGF₁₂₁, or one which contains the insertion in VEGF₁₆₆. Analysis of genomic clones in the area of putative mRNA splicing also shows an intron/exon structure consistent with alternative splicing.

Analysis of the nucleotide sequence of the VEGF gene indicates that VEGF is a member of the platelet-derived growth factor (PDGF) family. The amino acid sequence of VEGF exhibits approximately 20% homology to the sequences of the A and B chains of PDGF, as well as complete conservation of the eight cysteine residues found in both mature PDGF chains. VEGF also contains eight additional cysteine residues within the carboxy-terminal region. The amino-terminal sequence of VEGF is preceded by 26 amino acids corresponding to a typical signal sequence. The mature protein is generated directly following signal sequence cleavage without any intervening prosequence. The existence of a potential glycosylation site at Asn⁷⁴ is consistent with other evidence that VEGF is a glycoprotein, but the polypeptide has been reported to exist in both glycosylated and deglycosylated species.

Like other cytokines, VEGF can have diverse effects that depend on the specific biological context in which it is found. VEGF is a potent endothelial cell mitogen and directly contributes to induction of angiogenesis *in vivo* by promoting endothelial cell growth during normal development or during wound healing. A most striking property of VEGF is its specificity. It is mitogenic *in vitro* at 1 ng/ml for capillary and human umbilical vein endothelial cells, but not for adrenal cortex cells, corneal or lens epithelial cells, vascular smooth muscle cells, corneal endothelial cells, granulosa cells, keratinocytes, BHK-21 fibroblasts, 3T3 cells, rat embryo fibroblasts, human placental

fibroblasts and human sarcoma cells. The target cell specificity of VEGF is thus restricted to vascular endothelial cells. VEGF can trigger the entire sequence of events leading to angiogenesis and stimulates angiogenesis *in vivo* in the cornea and in a healing bone graft model. It is able to stimulate the proliferation of endothelial cells isolated from both small and large vessels. Expression of VEGF mRNA is temporally and spatially related to the physiological proliferation of capillary blood vessels in the ovarian corpus luteum or in the developing brain. VEGF expression is triggered by hypoxemia so that endothelial cell proliferation and angiogenesis appear to be especially stimulated in ischemic areas. VEGF is also a potent chemoattractant for monocytes. In addition, VEGF induces plasminogen activator and plasminogen activator inhibitor in endothelial cells.

Tumor cells release angiogenic molecules such as VEGF, and monoclonal antibodies to VEGF have been shown to inhibit the growth of certain types of tumor such as rhabdomyosarcoma. See Kim et al., "Inhibition of Vascular Endothelial Growth Factor-Induced Angiogenesis Suppresses Tumor Growth *in vivo*," *Nature*, 362:841-844 (1993). This suggests that blocking VEGF action is of potential therapeutic significance in treating tumors in general, and highly-vascularized, aggressive tumors in particular.

Summary of the Invention

It is an object of the invention to provide a new growth factor having the property of promoting proliferation of endothelial cells.

Another object of the invention is to provide isolated DNA sequences which encode a new growth factor which promotes proliferation of endothelial cells.

It is also an object of the invention to provide new products which may be useful in diagnostic and/or therapeutic applications.

These and other objects are achieved in accordance with the present invention by providing an isolated DNA sequence which codes for a protein having the property of promoting proliferation of endothelial cells or mesodermal cells, the
5 DNA sequence hybridizing under stringent conditions with a coding portion of the DNA sequence of Figure 1 or Figure 2.

In accordance with further aspects of the invention, the objects are also achieved by providing a mammalian protein having the property of promoting proliferation of
10 endothelial cells, which protein comprises an amino acid sequence substantially corresponding to the amino acid sequence of Figure 1 or the amino acid sequence of Figure 2, and by providing pharmaceutical preparations which comprise such proteins and antibodies which react with such proteins.

15 Clinical applications of the invention include diagnostic applications, acceleration of angiogenesis in wound healing, and inhibition of angiogenesis. Quantitation of VEGF-B in cancer biopsy specimens may be useful as an indicator of future metastatic risk. Topical application of
20 VEGF-B preparations to chronic wounds may accelerate angiogenesis and wound healing.

Brief Description of the Drawings

Figure 1 shows the nucleotide sequence of the (partial) cDNA clone of VEGF-B and the amino acid sequence of the
25 protein segment coded by the first reading frame of the cDNA;

Figure 2 repeats the nucleotide sequence of the (partial) cDNA clone of VEGF-B and the amino acid sequence
30 of the protein segment coded by the second reading frame of the cDNA; and

Figure 3 shows a comparison of the amino acid sequences of PDGF-A, PDGF-B, PlGF, VEGF and VEGF-B.

Detailed Description of Preferred Embodiments

The present invention thus is directed to a new vascular endothelial growth factor, hereinafter referred to as VEGF-B, which shares the angiogenic and other properties of VEGF, but which is distributed and expressed in tissues differently from VEGF.

VEGF-B is a member of the family of platelet derived growth factors and is a growth factor which promotes mitosis and proliferation of vascular endothelial cells and/or mesodermal cells. It is produced by expression of a DNA sequence which is hybridizable under stringent conditions with the DNA sequence depicted in Figure 1. Suitable hybridization conditions include, for example, 50% formamide, 5 x SSPE buffer, 5 x Denhardt's solution, 0.5% SDS and 100 µg/ml of salmon sperm DNA at 42°C overnight, followed by washing 2 x 30 minutes in 2 x SSC at 55°C.

The invention is also directed to an isolated and/or purified DNA which hybridizes under stringent conditions with the DNA sequence of Figure 1 or Figure 2. It is intended to include within the scope of the invention all angiogenic proteins encoded by DNA sequences which hybridize under stringent conditions to the DNA sequence of Figures 1 and 2.

In a further aspect, the invention is directed to antibodies of VEGF-B, and particularly to monoclonal antibodies.

A cDNA clone encoding murine VEGF-B was identified as follows. A cDNA library (E 14.5) derived from poly A⁺ mRNA isolated from 14.5 day old mouse embryos [Chevray P. and Nathans D., "Protein interaction cloning in yeast: identification of mammalian proteins that react with the leucine zipper of Jun," *Proc. Natl. Acad. Sci. USA*, 89:5789-93, 1992)] was screened for cellular proteins which potentially might interact with cellular retinoic acid-binding protein type 1 (CRABP-I) using a yeast two-hybrid interaction trap screening technique as described by Gyuris

J., Golemis E., Chertkov H. and Brent R., "Cdi1, a Human G1 and S Phase Protein Phosphatase That Associates with Cdk2," *Cell*, 75:791-803 (1993). This screening technique involves a fusion protein that contains a binding domain and that is known to be transcriptionally inert (the "bait"); reporter genes that have no basal transcription and that are bound by the bait; and an expression library which encodes proteins expressed as chimeras and whose amino termini contain an activation domain and other useful moieties (the "prey").

5 The screened library was a plasmid library in the yeast expression vector pPC67 obtained from Dr. Pierre Chevray of the Johns Hopkins University, School of Medicine, 725 North Wolfe St., Baltimore, MD 21205. A positive cDNA clone (pcif-2) was recovered from the screening. The positive

10 clone was sequenced using well known, conventional techniques and found to encode a protein highly homologous to VEGF and the other members of the PDGF family of growth factors. The 890 base pair SalI-NotI insert in the plasmid pPC67 was cloned into pBluescript and fully sequenced using

15 T7 and T3 vector primers together with internal primers. The plasmid pBluescript is commercially available from Stratagene Inc., LaJolla, California. The cDNA insert was found to be 886 base pairs long and to encode two polypeptides in different reading frames which were

20 homologous to the N-terminal end and the C-terminal end, respectively, of VEGF. This novel growth factor is referred to hereinafter as VEGF-B. The clone is partial and lacks approximately seven amino acids in the amino terminal region and the entire signal sequence of approximately twenty-eight

25 amino acids.

The protein is believed to interact with protein tyrosine kinase growth factor receptors. Details of such receptors are known in the art [See e.g. Wilks, A.F., "Protein Tyrosine Kinase Growth Factor Receptors and Their

30 Ligands in Development, Differentiation, and Cancer," *Adv. Cancer Res.*, 60:43-73 (1993)].

Various adult mouse tissues were tested for expression of transcripts corresponding to VEGF-B by Northern blotting. The size of the mRNA was 1.3-1.4 kb. A mouse multiple tissue Northern blot (MTN, Clontech) was probed with the 0.89 kb Sall-NotI fragment derived from the pPC67 yeast expression vectors described above. The probe was labelled with ³²P-dCTP using random priming (specific activity 10⁸-10⁹ cpm/μg of DNA). The blot was hybridized overnight at 42°C using 50% formamide, 5 x SSPE buffer, 2% SDS, 10 x Denhardt's solution, 100 μg/ml salmon sperm DNA and 1x10⁶ cpm of the labelled probe/ml. The blot was washed at room temperature for 2 x 30 min in 2 x SSC containing 0.05% SDS and then for 2 x 20 min at 52°C in 0.1 x SSC containing 0.1% SDS. The blot was then exposed at -70°C for three days using intensifying screens. Kodak XAR film was used. The relative expression levels as determined by visual examinations of the film are listed in the following table:

Table 1
Distribution of VEGF-B Transcripts in the Adult Mouse

Tissue	Relative Expression Level
Heart	+++++
Brain	+++
Spleen	(+)
Lung	++
Liver	+
Skeletal Muscle	++++
Kidney	+++
Testis	(+)

..... = very strong expression; .. = rather weak expression;
 = strong expression; . = weak expression;
 ... = moderate expression; ... = very weak expression.

A human multiple tissue Northern blot (MNT) from Clontech was probed using the murine partial cDNA to determine relative VEGF-B expression levels in various human tissues. The size of the transcript was 1.3-1.4 kb. The conditions were identical to those used for the mouse Northern blot described above. The relative VEGF-B transcript levels for the human Northern blot are listed in the following Table 2. For comparison purposes, Table 2 also lists relative expression level data from the literature for VEGF in various mammalian systems.

Table 2

Tissues	Relative Expression Levels			
	VEGF-B (Northern blot)	VEGF (from literature)		
		human	murine	guinea pig
heart	+++++	++	+++	+++
brain	+		+	+
placenta	+			
lung	+	++++		++
liver	(+)	++	(+)	+
skeletal muscle	++++		+++	+
kidney	+	++	+	++
pancreas	+++			
spleen	++		-	+
thymus	+		-	
prostate	+++			
testis	++			(+)
ovary	+++			-
small intestine	++			
colon	+++			
peripheral blood monocytes	+			

From a comparison of Table 1 and Table 2 it can be seen that mouse and human tissue expression levels of VEGF-B transcripts are relatively similar with the highest expression levels being found in heart and skeletal muscle. Significant differences may be seen in brain and kidney tissue. It should also be noted that tissues containing a large proportion of epithelial cells, such as prostate, pancreas and colon from which some of the most common human tumors originate, express relatively high levels of VEGF-B.

A comparison of the relative expression levels of VEGF and VEGF-B in human tissues shows some striking differences. VEGF is expressed rather weakly by human heart tissue, but VEGF-B is very strongly expressed by the same tissue. On the other hand, VEGF is strongly expressed by human lung tissue, but VEGF-B is only weakly expressed by human lung tissue. In a similar vein, human liver tissue expresses VEGF at a moderate level, but VEGF-B is expressed only very weakly. These data evidence that despite their general similarities, the actions of VEGF and VEGF-B are not completely identical.

Figure 1 shows the nucleotide sequence (SEQ ID NO:1) of the partial cDNA clone of VEGF-B and the amino acid sequence (SEQ ID NO:2) encoded in the first reading frame thereof. The DNA sequence of Figure 1 was obtained by conventional sequencing of a clone (pcif-2) in the yeast expression vector pPC67. The clone comprised 886 base pairs and encoded a part of murine VEGF-B.

The isolated cDNA sequence will hybridize with the mammalian genomic DNA, e.g. either murine or human, which contains the VEGF-B gene. In addition to the coding sequence, the genomic DNA will contain one or more promoter sequence(s) which give and direct expression of VEGF-B in one or more specific tissues. Thus the coding sequence of VEGF-B may be linked to an endothelial specific promoter which is specific to a certain type or types of tissue.

The full length protein is estimated to be approximately 120-125 amino acids in length.

The nucleotide sequence is translated in two different reading frames into two different amino acid sequences.

5 There is a stop codon (TGA) within the coding sequence at base pairs 309-311. Thus, VEGF-B comes in several splicing variants. The 5' end of the cloned cDNA sequence encodes an 102 amino acid long protein with significant homology to the N-terminal domains of VEGF, PlGF and PDGF A and B. In
10 particular, a number of cysteine residues are perfectly conserved within this group of proteins. In addition to the nucleotide sequence (SEQ ID NO:1), Figure 1 further depicts the deduced amino acid sequence (SEQ ID NO:2) of this first protein, which is 102 amino acids in length.

15 Translation of the C-terminal end of the cDNA (base pairs 308-475) in a different reading frame results in a protein which is highly homologous to the C-terminal part of VEGF..... Figure 2 again shows the nucleotide sequence (SEQ ID NO:1) of Figure 1, but this time includes the deduced
20 amino acid sequence (SEQ ID NO:3) of the second protein, which is encoded in the second reading frame and is 54 amino acids long. It thus appears that the VEGF-B gene encodes different proteins using alternative splicing of the primary transcript. The last part of the clone, encoding the second
25 peptide might be expressed as a functional protein in other spliced variants of VEGF-B.

Figure 3 shows a comparison of the amino acid sequence alignments of Platelet Derived Growth Factor A (PDGF-A), Platelet Derived Growth Factor B, (PDGF-B), Placenta Growth
30 Factor PlGF, Vascular Endothelial Growth Factor (VEGF) and the novel Vascular Endothelial Growth Factor B of the present invention (VEGF-B). As can be seen from this figure, the homologous relationship of the sequences is apparent, and VEGF-B is a structural homolog of the other
35 growth factors of this group. The boxes in Figure 3

indicate conserved cysteine residues in the respective protein amino acid sequences.

5 The afordescribed proteins may exist in combined association with an additional N-terminal sequence of approximately five (5) to ten (10) amino acids, as well as a further leader sequence of approximately twenty-eight (28) amino acids. Inasmuch such combined amino acid sequences exhibit the property of promoting the proliferation of endothelial cells and the DNA sequences which code for such
10 combined peptide sequences will hybridize under stringent conditions with the DNA sequence of Figures 1 and 2, such amino acid sequences and the DNA which codes for them are expressly contemplated to be within the scope of the present invention.

15 VEGF-B is synthesized normally in the endoplasmic reticulum of the source cell for subsequent export. Recombinant VEGF-B may be produced by inserting a DNA sequence encoding the VEGF-B protein together with a suitable operatively linked promoter and control sequences
20 into a suitable vector, such as the well known plasmid pBR322 or a derivative thereof, transforming or transfecting a suitable host cell, such as a Cos cell, with the resulting vector or other systems well known in the art and screening the resulting transformants for VEGF-B expression, and then
25 culturing cell lines which are positive for the expression of VEGF-B. Either a eukaryotic vector or a prokaryotic vector may be used, depending on the type of cell which is to be transfected or transformed therewith.

30 VEGF-B can be used as a growth factor for populations of endothelial cells in vitro. VEGF-B may be used to promote desirable angiogenesis, i.e. the formation of new blood vessels and capillaries. For example, it may be useful in promoting the development of the corpus luteum and endometrium as an aid to initiating and/or maintaining
35 pregnancy. Administration of VEGF-B may also be useful in supporting embryogenesis, as well as somatic growth and

vascular development and differentiation. Topical application of VEGF-B to wounds may be useful in promoting wound healing, and oral administration of VEGF-B may be useful to accelerate the healing of gastric and/or duodenal ulcers.

VEGF-B may exert proliferative effects on mesodermal cells either directly or via improvements in the blood supply.

Tumor assays for VEGF-B may be useful as indicators of metastatic risk. Assays of VEGF-B in body fluids or the tumor itself by histochemistry may be useful as a tumor prognostic factor. Furthermore, because tumor growth requires angiogenesis, administration of VEGF-B may also be useful in promoting tumor growth in laboratory animals in order to test anti-tumorigenic drugs. VEGF-B may also be useful to increase the microvasculature of hypoxic areas of tumors and make them more sensitive to radiation, radiation sensitizing drugs, etc.

The angiogenic action of VEGF-B may be useful in treating ischemic conditions. VEGF-B or agonists could be used to stimulate the development of collateral circulation in cases of arterial and/or venous obstruction, e.g. myocardial infarcts, ischaemic limbs, deep venous thrombosis, and/or postpartum vascular problems.

A VEGF-B/VEGF-B receptor system may be used as an assay system to detect small molecules as agonists/antagonists for development as new drugs.

Pharmaceutical compositions may be produced by admixing a pharmaceutically effective amount of VEGF-B protein with one or more suitable carriers or adjuvants such as water, mineral oil, polyethylene glycol, starch, talcum, lactose, thickeners, stabilizers, suspending agents, etc. Such compositions may be in the form of solutions, suspensions, tablets, capsules, creams, salves, ointments, or other conventional forms.

VEGF-B protein also can be used to produce antibodies. Such antibodies may be produced using conventional antibody production techniques. For example, specific monoclonal antibodies may be produced via immunization of fusion proteins obtained by recombinant DNA expression. Labelled monoclonal antibodies, in particular, should be useful in screening for conditions associated with abnormal levels of VEGF-B in the body. For example, assays of VEGF-B levels in blood or urine may be useful as a tumor marker. These monoclonal antibodies to VEGF-B also may be useful in inhibiting angiogenesis associated with high levels of VEGF-B in the body, e.g. in rapidly proliferating, angiogenesis-dependent tumors in mammals, and thereby may retard the growth of such tumors. Treatment may be effected, e.g., by twice weekly intraperitoneal injection of 10 to 500 μ g, preferably 50-100 μ g of monoclonal antibody. For the therapy of humans, chimerization or humanization of such monoclonal antibodies is to be preferred.

VEGF-B antagonists such as antibodies may be useful to inhibit new blood vessels in diabetic retinopathy, psoriasis, arthropathies and/or vascular tumors such as haemangiomas.

The foregoing description and examples have been set forth merely to illustrate the invention and are not intended to be limiting. Since modifications of the disclosed embodiments incorporating the spirit and substance of the invention may occur to persons skilled in the art, the invention should be construed to include everything within the scope of the appended claims and equivalents thereof.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Eriksson, Ulf
Olofsson, Birgitta
- (ii) TITLE OF INVENTION: VASCULAR ENDOTHELIAL GROWTH FACTOR-B AND DNA CODING THEREFOR
- (iii) NUMBER OF SEQUENCES: 3
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Evenson, McKeown, Edwards & Lenahan
 - (B) STREET: 1200 G Street, N.W., Suite 700
 - (C) CITY: Washington
 - (D) STATE: DC
 - (F) ZIP: 20005
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: 01-MAR-1995
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Evans, Joseph D
 - (B) REGISTRATION NUMBER: 26,269
 - (C) REFERENCE/DOCKET NUMBER: 1064/41979
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (202) 628-8800
 - (B) TELEFAX: (202) 628-8844

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 886 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CGGGACGCCC AGTGGTGCCA TGGATAGACG TTTATGCACG TGCCACATGC CAGCCCCAGGG	60
AGGTGGTGGT GCCTCTGAGC ATGGAACCTA TGGCAATGT GGTCAAACAA CTAGTCCCA	120
GCTGTGTGAC TTGTCAGCGC TTGTGTGGCT GCTGCCCTGA CGATGGCCTG GAATGTGTGC	180
CCACTGGGCA ACACCAAGTC CGAATGCAGA TCCTCATGAT CCAGTACCCG AGCAGTCAGC	240
TGGGGGAGAT GTCCCTGGAA GAACACAGCC AATGTGAATG CAGACCAAAA AAAAAAAGGA	300

GAGTGCTGTG AAGCCAGACA GCCCCAGGAT CCTCTGCCCG CCTTGCACCC AGCGCCGTCA 360
 ACGCCCTGAC CCCCAGACTT GCCCGTGCCG CTGCAGACGC CGCCGCTTCC TCCATTGCCA 420
 AGGGCGGGGC TTAGAGCTCA ACCCAGACAC CTGTAGGTGC CGGAAGCCGC GAAAGTGACA 480
 AGCTGCTTTC CAGACTCCAC TGGCCCGGCT GCTTTTATGG CCCTGCTTCA CAGGGACGAA 540
 GAGTGGAGCA CAGGCAAACC TCCTCAGTCT GGGAGGTCAC TGCCCCAGGA CCTGGACCTT 600
 TTAGAGAGCT CTCTCGCCAT CTTTTATCTC CCAGAGCTGC CATCTAACAA TTGTCAAGGA 660
 ACCTCATGTC TCACCTCAGG GGCCAGGGTA CTCTCTCACT TAACCACCCT GGTCAAGTGA 720
 GCATCTTCTG GCTGGCTGTC TCCCTCACT ATGAAAACCC CAACTTCTA CCAATAACGG 780
 GATTGGGGTT CTGTTATGAT AACTGTGACA CACACACACA CTCACACTCT GATAAAAGAG 840
 AACTCTGATA AAAGAGATGG AAGACACTAA AAAAAAAAAA AAAAAA 886

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 102 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Gly Arg Pro Val Val Pro Trp Ile Asp Val Tyr Ala Arg Ala Thr Cys
 1 5 10 15
 Gln Pro Arg Glu Val Val Val Pro Leu Ser Met Glu Leu Met Gly Asn
 20 25 30
 Val Val Lys Gln Leu Val Pro Ser Cys Val Thr Val Gln Arg Cys Gly
 35 40 45
 Gly Cys Cys Pro Asp Asp Gly Leu Glu Cys Val Pro Thr Gly Gln His
 50 55 60
 Gln Val Arg Met Gln Ile Leu Met Ile Gln Tyr Pro Ser Ser Gln Leu
 65 70 75 80
 Gly Glu Met Ser Leu Glu Glu His Ser Gln Cys Glu Cys Arg Pro Lys
 85 90 95
 Lys Lys Arg Arg Val Leu
 100

1. INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 55 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Lys	Pro	Asp	Ser	Pro	Arg	Ile	Leu	Cys	Pro	Pro	Cys	Thr	Gln	Arg	Arg
1				5				10					15		
Gln	Arg	Pro	Asp	Pro	Arg	Thr	Cys	Arg	Cys	Arg	Cys	Arg	Arg	Arg	Arg
		20					25					30			
Phe	Leu	His	Cys	Gln	Gly	Arg	Gly	Leu	Glu	Leu	Asn	Pro	Asp	Thr	Cys
		35					40					45			
Arg	Cys	Arg	Lys	Pro	Arg	Lys									
50						55									

What is claimed is:

1. An isolated DNA sequence which codes for a protein having the property of promoting proliferation of endothelial cells or mesodermal cells, said DNA sequence hybridizing under stringent conditions with a coding portion of the DNA sequence (SEQ ID NO:1) of Figure 1 or Figure 2.
2. A DNA sequence according to claim 1, wherein said DNA sequence is a cDNA sequence.
3. A DNA sequence according to claim 1, comprising a cDNA sequence corresponding to the DNA sequence of Figure 1 or Figure 2.
4. A DNA sequence according to claim 1, wherein said DNA sequence is a mammalian DNA sequence.
5. A DNA sequence according to claim 4, wherein said DNA sequence is a murine DNA sequence.
6. A DNA sequence according to claim 1, wherein said DNA sequence codes for a protein which promotes proliferation of vascular endothelial cells.
7. A vector comprising a DNA sequence according to claim 1.
8. A vector according to claim 7, wherein said vector is a eukaryotic vector.
9. A vector according to claim 7, wherein said vector is a prokaryotic vector.
10. A vector according to claim 7, wherein said vector is a plasmid.

11. A protein having the property of promoting proliferation of endothelial cells or mesodermal cells, said protein comprising an amino acid sequence substantially corresponding to the amino acid sequence of Figure 1 or the amino acid sequence of Figure 2.

12. A protein according to claim 11, wherein said protein comprises an amino acid sequence corresponding to the amino acid sequence (SEQ ID NO:2) of Figure 1.

13. A protein according to claim 11, wherein said protein comprises an amino acid sequence corresponding to the amino acid sequence (SEQ ID NO:3) of Figure 2.

14. A protein according to claim 11, wherein said protein is a mammalian protein.

15. A protein according to claim 14, wherein said protein is a murine protein.

16. A protein according to claim 11, wherein said protein promotes proliferation of vascular endothelial cells.

17. A pharmaceutical composition comprising an effective endothelial or mesodermal cell proliferation promoting amount of a protein according to claim 11, and at least one conventional pharmaceutical carrier or diluent.

18. An antibody which reacts with a protein according to claim 11.

19. An antibody according to claim 18, wherein said antibody is a monoclonal antibody.

20. A host cell transformed or transfected with a vector according to claim 7, such that said host cell expresses a protein having the property of promoting proliferation of endothelial or mesodermal cells.

21. A transformed host cell according to claim 20, wherein said host cell is a eukaryotic cell.

22. A cell according to claim 21, wherein said host cell is a COS cell.

23. A transformed host cell according to claim 20, wherein said host cell is a prokaryotic cell.

Abstract of the Disclosure

Polypeptide, VEGF-B, from the PDGF family of growth factors having the property of promoting mitosis and proliferation of vascular endothelial cells, DNA sequences encoding these polypeptides, pharmaceutical compositions containing them and antibodies which react with them. The VEGF-B polypeptides are useful in stimulating angiogenesis as well as in diagnostic applications.

Attorney Docket No. 1064/41979

DECLARATION AND POWER OF ATTORNEY - PATENT APPLICATION

As a below named inventor, I hereby declare that my citizenship, postal address and residence are as stated below; that I verily believe I am the original, first and sole inventor (if only one inventor is named below) or a joint inventor (if plural inventors are named below) of the invention entitled:

VASCULAR ENDOTHELIAL GROWTH FACTOR-B AND DNA CODING THEREFOR

the specification of which

is attached hereto, or

X was filed on March 1, 1995 as Application Serial No. 08/397,651 and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose all information known to be material to patentability as defined in 37 CFR §1.56. I hereby claim foreign priority benefits under Title 35, United States Code §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)

Priority Claimed

(Number)	(Country)	(Day/Month/Year)
(Number)	(Country)	(Day/Month/Year)

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose all information known to be material to patentability as defined in 37 CFR §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Application Serial No.:

(Filing Date)

(Status)

I hereby appoint as principal attorneys Herbert I. Cantor, Reg. No. 24,392; James F. McKernan, Reg. No. 25,406; Donald D. Evenson, Reg. No. 26,160; Joseph D. Evans, Reg. No. 26,269; Gary P. Edwards, Reg. No. 31,824, and Jeffrey D. Sanok, Reg. No. 32,169, to prosecute and transact all business in the Patent and Trademark Office connected with this application and any related United States and international applications. Please direct all communications to:

Evenson, McKernan, Edwards & Lenahan
1200 G Street, N.W., Suite 700
Washington, D.C. 20005
Telephone: (202) 628-8800
Facsimile: (202) 628-8844

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under §1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

INVENTOR: Ulf ERIKSSON

Citizenship: Sweden

Address/Residence: Bålsta, Sweden

May 2, 1995
Date

[Signature]
Signature

DECLARATION AND POWER OF ATTORNEY
Page 2

Attorney Docket No. 1064/41979

2-0
INVENTOR: Birgitta OLOFSSON

Citizenship: Sweden

Postal Address/Residence: Sundbyberg, Sweden

SEX

May 2, 1975
Date

Birgitta Olofsson
Signature

Reduced amino acid sequence of VEGF-B (Seq.1 and Seq.2)

09 397651

1	Gly	Arg	Pro	Val	Val	Pro	Trp	Ile	Asp	Val	Tyr	Ala	Arg	Ala	Thr	Cys	Gln	CG	17						
3	GGA	CGC	CCA	GTG	GTG	CCA	TGG	ATA	GAC	GTT	TAI	GCA	CGT	GCC	ACA	TGC	CAG		34						
	Pro	Arg	Glu	Val	Val	Val	Pro	Leu	Ser	Met	Glu	Leu	Met	Gly	Asn	Val	Val								
55	CCC	AGG	GAG	GTG	GTG	GTG	CGT	CTG	AGC	ATG	GAA	CTC	ATG	GGC	AAT	GTG	GTG		51						
	Lys	Gln	Leu	Val	Pro	Ser	Cys	Val	Thr	Val	Gln	Arg	Cys	Gly	Gly	Cys	Cys								
106	AAA	CAA	CTA	GTG	CCC	AGC	TGT	GTG	ACT	GTG	CAG	CGC	TGT	GGT	GGC	TGC	TGC		68						
	Pro	Asp	Asp	Gly	Leu	Glu	Cys	Val	Pro	Thr	Gly	Gln	His	Gln	Val	Arg	Met								
157	CCT	GAC	GAT	GGC	CTG	GAA	TGT	GTG	CCC	ACT	GGG	CAA	CAC	CAA	CTC	CGA	ATG		85						
	Gln	Ile	Leu	Met	Ile	Gln	Tyr	Pro	Ser	Ser	Gln	Leu	Gly	Glu	Met	Ser	Leu								
208	CAG	ATC	CTC	ATG	ATC	CAG	TAC	CCG	AGC	AGT	CAG	CTG	GGG	GAG	ATG	TCC	CTG		102						
	Glu	Glu	His	Ser	Gln	Cys	Glu	Cys	Arg	Pro	Lys	Lys	Lys	Arg	Arg	Val	Leu								
259	GAA	GAA	CAC	ACC	CAA	TGT	GAA	TGC	AGA	CCA	AAA	AAA	AAA	AGG	AGA	GTG	CTG		102						
	Stop																								
310	TGA	AGCC	CAG	ACG	CCCC	CAG	ATC	CTC	TG	CCCC	CGCT	TGC	AC	CC	CAG	CGG	CGT	CA	AC	CG	CC	TG	AC	CCCC	
376	GGAC	CTG	CCG	CTG	CCG	CTG	CAG	CGC	CGC	CGT	TCT	CCA	TG	CA	AG	GGG	GGG	GCT	TAG	AG	CT	CAA			
443	CCC	CAG	CAC	CTG	TAG	GTG	CCG	GAA	CGC	CGG	AA	AGT	GAC	AA	GCT	GCT	TTT	CC	AG	ACT	CC	AC	GGG	CCGG	
510	CTG	CTT	TTA	TGG	CCCT	GCT	TC	CA	GGG	AC	GA	AGT	GG	AG	CAC	AGG	CA	AA	CTC	CT	CAG	TC	GGG	AG	
577	GT	CAC	TG	CCCC	CAG	CGC	CGG	AC	CTT	TTA	GA	AG	CT	CT	CG	CA	TCT	TT	AT	CT	CC	CAG	AG	CT	CC
644	TCT	ACA	AAT	TG	CA	AG	CA	CT	CA	TG	TCT	CA	CT	CA	GGG	CC	AG	GGT	ACT	CT	CT	CAC	T	TA	ACC
711	TGG	CA	AGT	GA	CA	AT	CT	CT	GG	TGG	CT	CT	CC	CT	CA	TAT	GA	AA	AC	CC	AA	CT	CT	TA	CCA
778	ACG	GA	TT	TGG	TT	CT	CT	TT	AT	GA	TA	CT	GT	GA	CA	CA	CA	CA	CA	CA	CA	CA	CA	CA	CA
845	TCT	GAT	AAA	AG	AG	AT	GG	AA	TAC	ACT	AAAA	AAAA	AAAA	AAAA	AAAA	AAAA	AAAA	AAAA	AAAA	AAAA	AAAA	AAAA	AAAA	AAAA	

Figure 1

Reduced amino acid sequence of VEGF-B (Seq.1 and Seq.3)

09 397651

CGGGACGGC
10 CAGTGGTGGCATGCAATAGAGGTTTATGCACGTGCCACATGCCAGCCAGGGAGGTGGTGGCTCT
77 GACCATGGAACTCATGGGCAATGGTGTCAAACTAGTGGCCAGCTGTGTGACTGTGGAGCGCTGT
144 GGTGGCTGGTGGCTGAGGATGGGCTGGAAATGTGTGCCCACTGGGCAACCAAGTCCGAATGCAGA
211 TCTTCATGATCCAGTACCCGAGCAGTCAAGTGGGGAGATGTCTTGGGAAGAACACAGGCAATGTGA
Val Lys Pro Asp Ser Pro Arg
276 ATG CAG ACC AAA AAA AAG GAG AGT GCT GTG AAG CCA GAC AGC CCC AGG 7
Ile Leu Cys Pro Pro Cys Thr Gln Arg Arg Gln Arg Pro Asp Pro Arg Thr 24
351 ATG CTC TGC CCG CCT TGC ACC CAG CGC CGT CAA CGC CCT GAC CCC CGG ACC 4
Cys Arg Cys Arg Cys Arg Arg Arg Arg Phe Leu His Cys Gln Gly Arg Gly
381 TGC CGC TGC CGC TGC AGA CGC CGC CGC TTC CTC CAT TGC CAA GGG CGG GGC 56
Leu Glu Leu Asn Pro Asp Thr Cys Arg Cys Arg Lys Pro Arg Lys Stop
432 TTA GAG CTC AAC CCA GAC ACC TGT AGG TGC CGG AAG CCG CGA AAG TGA CAA
453 GGTGCTTTCCAGACTCCAGGGGCGCGCTGCTTTATGGCCCTGCTTCACAGGGACGAAAGTGGAG
550 CACAGGGCAAACTCTCTAGTCTGGGAGGTCACTGCCCGAGGACCTGGACCTTTAGAGAGCTCTCTC
617 GCCATCTTTATCTCCAGAGCTGCCATCTAACAAATTGTCAAGGAACCTCATGTCTCACCTCAGGGG
684 CCAAGGTACTCTCTCACTTAACCACTGCTCAAGTGAGCATCTTCTGGCTGGCTGTCTCCCTCAC
751 TATGAAACCCCAACTTTACCAATAACGGGATTTGGGTTCTGTTATGATAACTGTGACACACACA
816 CACACTCAGACTCTGATAAAGAGAACTCTGATAAAAGAGATGGAAACACTAAAAA
865 AAG

Figure 2

Amino acid sequence alignments of PDGFs, PLGF
VEGF and the novel homologues VEGF-B

397651

1 50
PDGF-A .MRTWACLLL LGGCYLALAL AEEAETPREL IERLARSQIE SIRDLOQLLE
PDGF-B MNRDMA.LFL PLCCYLALVS AEGDPTEEL YEMLSQHSIR SFDDLOQLLM
PLGFMPVHRL
VEGF-AMNT
VEGF-B
(Seq2)

51 100
PDGF-A IDSVGAEDA. LETSL.RARG SHAINM7EK P2VP1RRKRS ISEAIPAVK
PDGF-B KDSVDEGAE LDLMNTRAMS GVELESSRG RR.SIGSLAA AEPAVIAEY
PLGF FPCFLQLLAG LALPAUPPQC WALSAENGSS EV...EVVPT QEVWGREYK
VEGF-A LLSWVWNTLA LLL.YLHAK WSOAAPTTEG EQKSEVTKT KGVYGRSKY
VEGF-BGEPVVEN IDVYARATG
(Seq2)

101 150
PDGF-A TRIVITEIFR SQVDPTISANT LIMPPTEVK RIGGCTSS VKDPSRVNH
PDGF-B TRIEVFOISR NLIDRTNANT LVWPPTEVQ RSGCKNRN VQNASQVCH
PLGF ALERLVQVVS EYPS..EVEN MFSPS3LL RIGGCGDEN LQVPVETAN
VEGF-A PIETLVDFIQ EYPD..EIEY IFKPS77LM RACGCTEA LQVPTSESN
VEGF-B PREVVVPLSM ELNG..NVVK QLVPS7TVQ RIGGCTDDG LQV7TAENH
(Seq2)

151 200
PDGF-A RSVKVARVEY VRKKPKLKEV QVRLEENLE AAT.....SNLNPZ
PDGF-B RSVQVVKIEI VWSKP1FKKA TVTLECHLAC KQETIVTPRP VTRSPGTSAS
PLGF VTNQLLKINS GDSFST...V ELTFSQHVFC KQ.....RPL REX...MKPE
VEGF-A ITHQDKRIKP WQSONT...E RMSFLQHSFC KQ.....RPK FOR...TKPE
VEGF-B VTNQILNIQY PSSQ..L...G RMSLEINSK KQ.....RPR KQRRV.....
(Seq2)

VEGF-BVKPD
(Seq2)

201 250
PDGF-A HREETDVR.
PDGF-B ORAKTPGARV TIRTVRIRAP PKGKMYFKH THDYAALKET LGA.....
PLGF KGGDAVPRR.
VEGF-A NMCEETERR KHLFVDDPQTNTCSAAQLEL NERT.....
VEGF-B
(Seq2)
VEGF-B SPRILPCTQRR QRPDPKTPRPE LHTGRGLEL NPDQ.....
(Seq2)

251
PDGF-A ..
PDGF-B ..
PLGF ..
VEGF-A RP
VEGF-B ..
(Seq2)
VEGF-B ..
(Seq3) RK

Figure 3